



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Biofeedback as a Treatment of Urinary Incontinence in Adults

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 173

BCBSA Reference Number: 2.01.27 (For Plans internal use only)

Related Policies

- Pelvic Floor Stimulation as a Treatment of Urinary Incontinence, [#470](#)
- Transvaginal and Transurethral Radiofrequency Tissue Remodeling for Urinary Stress Incontinence, [#523](#)
- Periurethral Bulking Agents for the Treatment of Incontinence, [#471](#)
- Sacral Nerve Neuromodulation/Stimulation for Pelvic Floor Dysfunction, [#153](#)
- Percutaneous Tibial Nerve Stimulation for Voiding Dysfunction, [#583](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Biofeedback in the outpatient setting as a treatment of urinary incontinence in adults is [INVESTIGATIONAL](#).

Unsupervised home use of biofeedback for treatment of urinary incontinence is [INVESTIGATIONAL](#).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

According to the policy statement above, the following CPT codes are considered investigational for the conditions listed for **Commercial Members: Managed Care (HMO and POS), and PPO, Indemnity:**

CPT Codes

CPT codes:	Code Description
90875	Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (eg, insight oriented, behavior modifying or supportive psychotherapy); approximately 20-30 minutes
90876	Individual psychophysiological therapy incorporating biofeedback training by any modality (face-to-face with the patient), with psychotherapy (eg, insight oriented, behavior modifying or supportive psychotherapy); approximately 45-50 minutes
90901	Biofeedback training by any modality
90912	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; initial 15 minutes of one-on-one physician or other qualified health care professional contact with the patient
90913	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; each additional 15 minutes of one-on-one physician or other qualified health care professional contact with the patient (List separately in addition to code for primary procedure)

The following CPT code is considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

HCPCS Codes

HCPCS codes:	Code Description
S9002	Intra-vaginal motion sensor system, provides biofeedback for pelvic floor muscle rehabilitation device

Description

Biofeedback

Biofeedback is intended to teach patients self-regulation of certain physiologic processes not normally considered to be under voluntary control. The technique involves feedback on a variety of types of information not commonly available to the patient, followed by a concerted effort on the part of the patient to use this feedback to help alter the physiologic process in some specific way. Biofeedback has been proposed as a treatment for a variety of diseases and disorders, including anxiety, headaches, hypertension, movement disorders, incontinence, pain, asthma, Raynaud disease, and insomnia. Biofeedback training is done either in individual or group sessions and as a single therapy or in combination with other therapies designed to teach relaxation. A typical program consists of 10 to 20 training sessions of 30 minutes each. Training sessions are performed in a quiet, nonarousing environment. Subjects are instructed to use mental techniques to affect the physiologic variable monitored, and feedback is provided for the successful alteration of the physiologic parameter. This feedback may be in the form of signals, such as lights or tone, verbal praise, or other auditory or visual stimuli.

Biofeedback, in conjunction with pelvic floor muscle training, is a possible treatment modality for stress, urge, mixed, and overflow urinary incontinence because it may enhance awareness of body functions and the learning of exercises to train pelvic muscles. Several proposed biofeedback methods may be employed to treat urinary incontinence, including vaginal cones or weights, perineometers, and electromyographic systems with vaginal and rectal sensors.

The various forms of biofeedback mainly differ in the nature of the disease or disorder under treatment, the biologic variable that the subject attempts to control, and the information that is fed back to the subject. Biofeedback techniques include peripheral skin temperature feedback, blood-volume-pulse feedback (vasoconstriction and dilation), vasoconstriction training (temporalis artery), and electromyographic biofeedback; they may be used alone or in conjunction with other therapies (eg, relaxation, behavioral management, medication).

Summary

Description

Biofeedback is a technique to teach patients self-regulation of physiologic processes not generally considered to be under voluntary control; a variety of approaches and devices are available. Biofeedback, in conjunction with pelvic floor muscle training, is proposed as a treatment of urinary incontinence.

Summary of Evidence

For individuals who have urinary incontinence (women) who receive biofeedback with pelvic floor muscle training (PFMT), the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, functional outcomes, and quality of life. A comparative effectiveness review did not find a statistically significant difference in continence rates when patients received PFMT with or without biofeedback. Other systematic reviews evaluating biofeedback and/or verbal feedback as part of treatment for urinary incontinence found improvement in some outcomes but not others. There is a lack of consistent evidence from well-designed trials that biofeedback effectively treats urinary incontinence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have post-prostatectomy urinary incontinence, the evidence includes an RCT and systematic reviews that compared PFMT with or without biofeedback. Relevant outcomes are symptoms, functional outcomes, and quality of life. Results of these data were mixed and did not consistently report significantly improved outcomes when biofeedback was added to the intervention. The timing and delivery of the intervention were not well-defined. Additional well-designed trials are needed that demonstrate the superiority of biofeedback with PFMT over PFMT alone. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who will undergo radical prostatectomy, the evidence includes RCTs that have evaluated the efficacy of biofeedback with PFMT compared with PFMT without biofeedback for prevention of prostatectomy-related urinary incontinence. These trials generally did not report improved outcomes with biofeedback added to the intervention. The timing and delivery of the intervention were not well-defined. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
4/2024	Clarified coding information.
10/2023	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
10/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
9/2021	Annual policy review. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
10/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2020	Clarified coding information.

10/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2017	New references added from BCBSA National medical policy.
11/2015	Clarified coding information.
8/2015	Annual policy review. New references added
9/2014	Annual policy review. New references added
5/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
10/2013	Annual policy review. New references added
5/2013	Annual policy review. New references added
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
9/2011	Reviewed - Medical Policy Group – Urology, Obstetrics and Gynecology. No changes to policy statements.
6/2011	Reviewed - Medical Policy Group – Orthopedics, Rehabilitation Medicine and Rheumatology. No changes to policy statements.
7/2010	Reviewed - Medical Policy Group – Orthopedics, Rehabilitation Medicine and Rheumatology. No changes to policy statements.
6/2010	Reviewed - Medical Policy Group - Urology. No changes to policy statements.
3/1/2010	Medical Policy 173 effective 3/1/2010 describing covered and non-covered indications.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Biofeedback. TEC Assessments. 1995;Volume 10:Tab 25.
2. Burgio KL, Goode PS, Locher JL, et al. Behavioral training with and without biofeedback in the treatment of urge incontinence in older women: a randomized controlled trial. JAMA. Nov 13 2002; 288(18): 2293-9. PMID 12425706
3. Muscle strength. Physiopedia. Accessed June 23,2023. https://www.physio-pedia.com/Muscle_Strength
4. Pelvic muscle function and strength. Physiopedia. Accessed June 23,2023. https://www.physio-pedia.com/Pelvic_Floor_Muscle_Function_and_Strength
5. Zhu D, Xia Z, Yang Z. Effectiveness of physiotherapy for lower urinary tract symptoms in postpartum women: systematic review and meta-analysis. Int Urogynecol J. Mar 2022; 33(3): 507-521. PMID 34302516
6. Wu X, Zheng X, Yi X, et al. Electromyographic Biofeedback for Stress Urinary Incontinence or Pelvic Floor Dysfunction in Women: A Systematic Review and Meta-Analysis. Adv Ther. Aug 2021; 38(8): 4163-4177. PMID 34176082
7. Mateus-Vasconcelos ECL, Ribeiro AM, Antônio FI, et al. Physiotherapy methods to facilitate pelvic floor muscle contraction: A systematic review. Physiother Theory Pract. Jun 2018; 34(6): 420-432. PMID 29278967
8. Laycock J, Jerwood D. Pelvic floor muscle assessment: the PERFECT scheme. Physiotherapy. 2001;87(12):631-642. Accessed June 23,2023. <https://www.sciencedirect.com/science/article/abs/pii/S003194060561108X>
9. Moroni RM, Magnani PS, Haddad JM, et al. Conservative Treatment of Stress Urinary Incontinence: A Systematic Review with Meta-analysis of Randomized Controlled Trials. Rev Bras Ginecol Obstet. Feb 2016; 38(2): 97-111. PMID 26883864
10. Shamliyan T, Wyman J, Kane RL, et al. Nonsurgical Treatments for Urinary Incontinence in Adult Women: Diagnosis and Comparative Effectiveness. Comparative Effectiveness Review No. 36. Rockville, MD: Agency for Healthcare Research and Quality; 2012.

11. Herderschee R, Hay-Smith EJ, Herbison GP, et al. Feedback or biofeedback to augment pelvic floor muscle training for urinary incontinence in women. *Cochrane Database Syst Rev.* Jul 06 2011; (7): CD009252. PMID 21735442
12. Hagen S, Elders A, Stratton S, et al. Effectiveness of pelvic floor muscle training with and without electromyographic biofeedback for urinary incontinence in women: multicentre randomised controlled trial. *BMJ.* Oct 14 2020; 371: m3719. PMID 33055247
13. Williams KS, Assassa RP, Gillies CL, et al. A randomized controlled trial of the effectiveness of pelvic floor therapies for urodynamic stress and mixed incontinence. *BJU Int.* Nov 2006; 98(5): 1043-50. PMID 17034605
14. Hirakawa T, Suzuki S, Kato K, et al. Randomized controlled trial of pelvic floor muscle training with or without biofeedback for urinary incontinence. *Int Urogynecol J.* Aug 2013; 24(8): 1347-54. PMID 23306768
15. Pereira VS, de Melo MV, Correia GN, et al. Vaginal cone for postmenopausal women with stress urinary incontinence: randomized, controlled trial. *Climacteric.* Feb 2012; 15(1): 45-51. PMID 22066898
16. Bertotto A, Schvartzman R, Uchôa S, et al. Effect of electromyographic biofeedback as an add-on to pelvic floor muscle exercises on neuromuscular outcomes and quality of life in postmenopausal women with stress urinary incontinence: A randomized controlled trial. *Neurourol Urodyn.* Nov 2017; 36(8): 2142-2147. PMID 28508398
17. Özlü A, Yıldız N, Öztekin Ö. Comparison of the efficacy of perineal and intravaginal biofeedback assisted pelvic floor muscle exercises in women with urodynamic stress urinary incontinence. *Neurourol Urodyn.* Nov 2017; 36(8): 2132-2141. PMID 28345778
18. Goode PS, Burgio KL, Johnson TM, et al. Behavioral therapy with or without biofeedback and pelvic floor electrical stimulation for persistent postprostatectomy incontinence: a randomized controlled trial. *JAMA.* Jan 12 2011; 305(2): 151-9. PMID 21224456
19. Hsu LF, Liao YM, Lai FC, et al. Beneficial effects of biofeedback-assisted pelvic floor muscle training in patients with urinary incontinence after radical prostatectomy: A systematic review and metaanalysis. *Int J Nurs Stud.* Aug 2016; 60: 99-111. PMID 27297372
20. Johnson EE, Mamoulakis C, Stoniute A, et al. Conservative interventions for managing urinary incontinence after prostate surgery. *Cochrane Database Syst Rev.* Apr 18 2023; 4(4): CD014799. PMID 37070660
21. MacDonald R, Fink HA, Huckabay C, et al. Pelvic floor muscle training to improve urinary incontinence after radical prostatectomy: a systematic review of effectiveness. *BJU Int.* Jul 2007; 100(1): 76-81. PMID 17433028
22. Oh JJ, Kim JK, Lee H, et al. Effect of personalized extracorporeal biofeedback device for pelvic floor muscle training on urinary incontinence after robot-assisted radical prostatectomy: A randomized controlled trial. *Neurourol Urodyn.* Feb 2020; 39(2): 674-681. PMID 31793032
23. Tienforti D, Sacco E, Marangi F, et al. Efficacy of an assisted low-intensity programme of perioperative pelvic floor muscle training in improving the recovery of continence after radical prostatectomy: a randomized controlled trial. *BJU Int.* Oct 2012; 110(7): 1004-10. PMID 22332815
24. Wille S, Sobottka A, Heidenreich A, et al. Pelvic floor exercises, electrical stimulation and biofeedback after radical prostatectomy: results of a prospective randomized trial. *J Urol.* Aug 2003; 170(2 Pt 1): 490-3. PMID 12853806
25. Bales GT, Gerber GS, Minor TX, et al. Effect of preoperative biofeedback/pelvic floor training on continence in men undergoing radical prostatectomy. *Urology.* Oct 01 2000; 56(4): 627-30. PMID 11018619
26. ACOG Practice Bulletin No. 155: Urinary Incontinence in Women. *Obstet Gynecol.* Nov 2015; 126(5): e66-e81. PMID 26488524
27. Kobashi KC, Albo ME, Dmochowski RR, et al. Surgical Treatment of Female Stress Urinary Incontinence: AUA/SUFU Guideline. *J Urol.* Oct 2017; 198(4): 875-883. PMID 28625508
28. Kobashi KC, Vasavada S, Bloschichak A, et al. Updates to Surgical Treatment of Female Stress Urinary Incontinence (SUI): AUA/SUFU Guideline (2023). *J Urol.* Jun 2023; 209(6): 1091-1098. PMID 37096580
29. Sandhu JS, Breyer BB, Comiter C, et al. Incontinence after Prostate Treatment: AUA/SUFU Guideline (2019). American Urological Association. 2019. Accessed June 23, 2023. <https://www.auanet.org/guidelines/incontinence-after-prostate-treatment>
30. National Institute for Health and Care Excellence (NICE) Guideline. Urinary Incontinence and Pelvic Organ Prolapse in Women: Management. NICE Guideline. 2019. Accessed June 23, 2023. <https://www.nice.org.uk/guidance/ng123>
31. Centers for Medicare & Medicaid Services. National coverage decision (NCD) for biofeedback therapy for the treatment of urinary incontinence (30.1.1). 2001. <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=42>. Accessed June 23, 2023.